

# A new chemodynamical tool to study the evolution of galaxies in the local Universe: a quick and accurate numerical technique to compute gas cooling rate for any chemical composition

Nicolas Champavert<sup>1</sup> and Hervé Wozniak<sup>1</sup>

Université Lyon 1, CRAL, Observatoire de Lyon, 9 avenue Charles André,  
Saint-Genis Laval cedex, F-69561, France [champavert@obs.univ-lyon1.fr](mailto:champavert@obs.univ-lyon1.fr)

**Summary.** We have developed a quick and accurate numerical tool to compute gas cooling whichever its chemical composition.

## 1 Introduction

Metal abundances in galaxies vary from place to place and with time. These variations reflect the star formation history within galaxies because the chemical enrichment results from synthesis of heavy elements by successive generations of stars. Gas radiative cooling is very sensitive to the ISM chemical composition. Cooling influences both the gas dynamics and the star formation rate. Therefore we need to follow self-consistently dynamical and chemical evolution. We have developed a method to compute in a very short CPU time accurate cooling functions dependent on the exact abundances of elements. It will be used in high resolution chemodynamical simulations of galaxies.

## 2 Description of the method

It is well known that cooling rates depend on gas chemical composition. A gas with solar composition has a cooling rate greater than without metals by more than an order of magnitude for some temperatures (Fig. 1). Cooling rates are generally interpolated between rates computed for different metallicities with solar abundance ratios (see [1] for instance) or with solar abundance ratios and enhanced abundances for some elements (see [2] for instance). The gas abundances are obviously not always solar because the gas ejected from stars by winds and SNe has a non-solar composition. Hence we need to take into account the real abundances of each element in the gas to build more realistic cooling functions whichever the chemical composition. However, computing cooling rates with Mappings III for each gas particle at each timestep of a




**Fig. 1.** Cooling rates as a function of temperature for different metallicities computed with Mappings III. The abundances used in calculations are taken from [2]

chemodynamical simulation is a very CPU time consuming task. We thus have developed a recipe to reconstruct the cooling curve on the fly.

We consider the ISM as an optically thin gas in collisional ionization equilibrium. Calculations are performed in the temperature range  $10^4$  K to  $10^8$  K using Mappings III, successor of Mappings II whose cooling computations details are described in [2]. All the abundances of elements are relative to hydrogen. The hydrogen density is  $1.0 \text{ cm}^{-3}$ , independent of temperature. We drop the temperature dependence of cooling rates ( $\Lambda(T)$ ) for the sake of legibility.

Many cooling processes depend on the density of electrons and on the abundances of elements. A simple hypothesis is to suppose that cooling rates are proportional to these two quantities. Thus, we can try to reconstruct the total cooling curve  $\Lambda_{tot}$  with a linear combination of individual curves for each element. We first need to compute these individual curves. We obtain  $\Lambda_H$  with a purely hydrogen gas. For all the heavier elements used for calculations in Mappings III (He, C, N, O, Ne, Na, Mg, Al, Si, S, Cl, Ar, Ca, Fe, Ni), we have calculated cooling rates for mixtures of hydrogen and one element  $X_i$  at its solar abundance  $\Lambda_{X_i,H}$ . Individual cooling curves  $\Lambda_{X_i}$  have then been obtained subtracting  $\Lambda_H$  from each  $\Lambda_{H,X_i}$ :  $\Lambda_{X_i} = \Lambda_{H,X_i} - \Lambda_H$ . For a given chemical composition, we then compute the quantity  $\alpha_{X_i}$  for all the elements previously listed, where  $\alpha_{X_i}$  is the abundance of element  $X_i$  normalized by its solar abundance:  $\alpha_{X_i} = n_{X_i}/n_{X_i\odot}$ . For helium, our hypothesis is too simple. Its cooling rate cannot be considered proportional to its abundance with a good approximation as there are too strong non-linearities. We choose to linearly interpolate in a grid of precomputed curves for different abundances to obtain the helium cooling curve  $\Lambda_{He,\alpha_{He}}$ . As hydrogen and helium are the two most abundant elements, we assume that they are the only sources of electrons. We indeed consider that the contribution of the metals are negligible because of their much lower abundances in the ISM. By construction, our individual cooling functions for metals take into account the electrons of the




**Fig. 2. (left panel)** Cooling rates for several elements at solar abundances. The curve for other elements is the sum of the contributions of N, Ne, Na, Mg, Al, Si, S, Cl, Ar, Ca and Ni. The solid broad line  $\Lambda_{tot}$  is the cooling curve for solar abundances reconstructed using our recipe. **(right panel)** Percentage of relative error made when reconstructing cooling functions with our recipe for different compositions

element and hydrogen. Thus we only need to add the contribution of electrons provided by the ionization of helium  $ne(He)$ . We add the terms for hydrogen, helium and metals to build  $\Lambda_{tot}$  so that we finally obtain the following approximation:  $\Lambda_{tot} \approx \Lambda_H + \Lambda_{He, \alpha_{He}} + (1 + ne(He)) \sum_{metals} \alpha_{X_i} \Lambda_{X_i}$

To check the accuracy of our tool, we have compared cooling curves computed with Mappings III and the ones reconstructed with our recipe for various chemical compositions. We choose a solar composition to test a well-known chemical composition. Carbon, oxygen and iron are three of the most important coolants (Fig. 2, left panel). Furthermore, they are three of the most abundant elements in stellar winds and supernovae ejecta. Thus, we increase by a factor of 10 the abundance of these three important coolants. This factor is somewhat arbitrary and was only chosen to test the accuracy of our recipe in the case of a very high-enhanced gas. In all cases, the relative errors remain below a few percent (Fig. 2, right panel). The errors due to the computation of cooling rates are thus comparable to the others coming from, for instance, the hydrodynamical scheme.

## References

1. Boehringer, H. and Hensler, G.: A&A **215**, 147 (1989)
2. Sutherland, R.S. and Dopita, M. A.: ApJS **88**, 253 (1993)